

DISTANCE-SQUARED MAPPINGS

SHUNSUKE ICHIKI AND TAKASHI NISHIMURA

ABSTRACT. A distance-squared function is one of the most significant functions in the application of singularity theory to differential geometry. In this paper, we define naturally extended mappings of distance-squared functions, wherein each component is a distance-squared function. We investigate the properties of these mappings from the viewpoint of differential topology.

1. INTRODUCTION

Let $\mathbb{R}^n$ be the n -dimensional Euclidean space; thus, a point $x \in \mathbb{R}^n$ is n -tuple $x = (x_1, \dots, x_n)$ of real numbers. Let $d : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ be the n -dimensional Euclidean distance,

$$d(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2},$$

where $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$. Let p be a given point in $\mathbb{R}^n$. Then, the mapping $d_p : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by $d_p(x) = d(p, x)$, which is called a *distance function*.

Definition 1.1. Let $p_1, \dots, p_\ell$ ($\ell \geq 1$) be given points in $\mathbb{R}^n$. Then, the following mapping $d_{(p_1, \dots, p_\ell)} : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ is called a *distance mapping*:

$$d_{(p_1, \dots, p_\ell)}(x) = (d(p_1, x), \dots, d(p_\ell, x)).$$

A distance mapping is one in which each component is a distance function.

Let S^n be the n -dimensional unit sphere in $\mathbb{R}^{n+1}$

Proposition 1.1. Let $i : S^1 \rightarrow i(S^1) \subset \mathbb{R}^2$ be a homeomorphism. Then, there exist two points $p_1, p_2 \in i(S^1)$ such that $d_{(p_1, p_2)} \circ i : S^1 \rightarrow \mathbb{R}^2$ is homeomorphic to the image $d_{(p_1, p_2)} \circ i(S^1)$.

Proposition 1.1 is proved in Section 4. Although Proposition 1.1 is applicable even if a mapping i is not differentiable anywhere, it appears difficult to derive higher-dimensional extensions of the proposition.

On the other hand, the differentiable version of higher-dimensional extensions can be obtained as follows. All manifolds and mappings in this paper belong to class C^∞ except Proposition 1.1.

Definition 1.2. Let $p_1, \dots, p_\ell$ ($\ell \geq 1$) be given points in $\mathbb{R}^n$. Then, the following mapping $D_{(p_1, \dots, p_\ell)} : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ is called a *distance-squared mapping*:

$$D_{(p_1, \dots, p_\ell)}(x) = (d^2(p_1, x), \dots, d^2(p_\ell, x)).$$

2010 *Mathematics Subject Classification.* 57R40, 57R42.

Key words and phrases. distance-squared mapping; singularity; embedding; immersion; normal crossing.




FIGURE 1. Figure of Proposition 1.1

Note that $D_{(p_1, \dots, p_\ell)}$ always has a singular point. Nevertheless, Theorems 1.1 and 1.2 hold as follows.

Theorem 1.1. *Let M be an m -dimensional closed manifold ($m \geq 1$), and let $i : M \rightarrow \mathbb{R}^\ell$ ($m+1 \leq \ell$) be an embedding. Then, there exist $p_1, \dots, p_{m+1} \in i(M)$, $p_{m+2}, \dots, p_\ell \in \mathbb{R}^\ell$ such that $D_{(p_1, \dots, p_\ell)} \circ i : M \rightarrow \mathbb{R}^\ell$ is an embedding.*

For the definition of embedding, see [1].

Corollary 1.1. *Let M be an m -dimensional closed manifold ($m \geq 1$), and let $i : M \rightarrow \mathbb{R}^{m+1}$ be an embedding. Then, there exist $p_1, \dots, p_{m+1} \in i(M)$ such that $D_{(p_1, \dots, p_{m+1})} \circ i : M \rightarrow \mathbb{R}^{m+1}$ is an embedding.*

Let M and N be manifolds. An immersion $f : M \rightarrow N$ (for the definition of immersion, see [1]) is said to be *with normal crossing at a point $y \in N$* if $f^{-1}(y)$ is a finite set $\{x_1, x_2, \dots, x_n\}$ and for any subsets $\{\lambda_1, \lambda_2, \dots, \lambda_s\} \subset \{1, 2, \dots, n\}$ ($s \leq n$),

$$\text{codim} \left(\bigcap_{j=1}^s df_{x_{\lambda_j}}(T_{x_{\lambda_j}} M) \right) = \sum_{j=1}^s \text{codim} (df_{x_{\lambda_j}}(T_{x_{\lambda_j}} M)),$$

where $\text{codim } H = \dim T_y N - \dim H$ for a linear subspace $H \subset T_y N$. An immersion $f : M \rightarrow N$ is said to be *with normal crossing* if f is a mapping with normal crossing at any point $y \in N$.

Theorem 1.2. *Let M be an m -dimensional closed manifold ($m \geq 1$), and let $i : M \rightarrow \mathbb{R}^\ell$ ($m+1 \leq \ell$) be an immersion with normal crossing. Then, there exist $p_1, \dots, p_{m+1} \in i(M)$, $p_{m+2}, \dots, p_\ell \in \mathbb{R}^\ell$ such that $D_{(p_1, \dots, p_\ell)} \circ i : M \rightarrow \mathbb{R}^\ell$ is an immersion with normal crossing.*

Corollary 1.2. *Let M be an m -dimensional closed manifold ($m \geq 1$), and let $i : M \rightarrow \mathbb{R}^{m+1}$ be an immersion with normal crossing. Then, there exist $p_1, \dots, p_{m+1} \in i(M)$ such that $D_{(p_1, \dots, p_{m+1})} \circ i : M \rightarrow \mathbb{R}^{m+1}$ is an immersion with normal crossing.*

L -points $p_1, \dots, p_\ell \in \mathbb{R}^n$ ($1 \leq \ell \leq n+1$) are said to be *in general position* if $\ell = 1$ or $\overrightarrow{p_1 p_2}, \dots, \overrightarrow{p_1 p_\ell}$ ($2 \leq \ell \leq n+1$) are linearly independent, where $\overrightarrow{p_i p_j}$ stands for $(p_{j1} - p_{i1}, \dots, p_{jn} - p_{in})$ ($p_i = (p_{i1}, \dots, p_{in})$, $p_j = (p_{j1}, \dots, p_{jn}) \in \mathbb{R}^n$). A mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ is said to be *$\mathcal{A}$ -equivalent* to a mapping $g : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ if there exist diffeomorphisms $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\psi : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ such that $\psi \circ f \circ \varphi = g$. A mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ ($\ell \leq n$) is called *the normal form of definite fold mappings* if $f(x_1, \dots, x_n) = (x_1, \dots, x_{\ell-1}, x_\ell^2 + \dots + x_n^2)$. The properties of distance-squared

mappings are of significance in the proofs of Theorems 1.1 and 1.2. It turns out that distance-squared mappings are $\mathcal{A}$ -equivalent to the normal form of definite fold mappings if ℓ -points $p_1, \dots, p_\ell \in \mathbb{R}^n$ ($2 \leq \ell \leq n$) are in general position (see Section 2).

In Section 2, a fundamental property of distance-squared mappings is shown. The proofs of Theorems 1.1 and 1.2 are given in Section 3. Finally, in Section 4, the proof of Proposition 1.1 is given for the sake of readers' convenience.

2. PRELIMINARIES

Proposition 2.1.

(I) Let ℓ, n be integers such that $2 \leq \ell \leq n$, and let $p_1, \dots, p_\ell \in \mathbb{R}^n$ be in general position. Then, $D_{(p_1, \dots, p_\ell)} : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ is $\mathcal{A}$ -equivalent to the normal form of definite fold mappings.

(II) Let ℓ, n be integers such that $2 \leq n < \ell$, and let $p_1, \dots, p_{n+1} \in \mathbb{R}^n$ be in general position. Then, $D_{(p_1, \dots, p_\ell)} : \mathbb{R}^n \rightarrow \mathbb{R}^\ell$ is $\mathcal{A}$ -equivalent to the inclusion $(x_1, \dots, x_n) \mapsto (x_1, \dots, x_n, 0, \dots, 0)$.

2.1. **Proof of (I).** Let $H_1 : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the following diffeomorphism.

$$\begin{aligned} & H_1(x_1, x_2, \dots, x_\ell) \\ &= \left(\frac{1}{2} \left(x_1 - x_2 + \sum_{j=1}^n (p_{1j} - p_{2j})^2 \right), \dots, \frac{1}{2} \left(x_1 - x_\ell + \sum_{j=1}^n (p_{1j} - p_{\ell j})^2 \right), x_1 \right) \end{aligned}$$

The composition of $D_{(p_1, \dots, p_\ell)}$ and H_1 is as follows.

$$\begin{aligned} & (H_1 \circ D_{(p_1, \dots, p_\ell)})(x_1, x_2, \dots, x_n) \\ &= \left(\sum_{j=1}^n (p_{2j} - p_{1j})(x_j - p_{1j}), \dots, \sum_{j=1}^n (p_{\ell j} - p_{1j})(x_j - p_{1j}), \sum_{j=1}^n (x_j - p_{1j})^2 \right) \end{aligned}$$

Let $H_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the diffeomorphism defined by

$$H_2(x_1, x_2, \dots, x_n) = (x_1 + p_{11}, x_2 + p_{12}, \dots, x_n + p_{1n}).$$

The composition of $H_1 \circ D_{(p_1, \dots, p_\ell)}$ and H_2 is as follows:

$$\begin{aligned} & (H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2)(x_1, x_2, \dots, x_n) \\ &= \left(\sum_{j=1}^n (p_{2j} - p_{1j})x_j, \dots, \sum_{j=1}^n (p_{\ell j} - p_{1j})x_j, \sum_{j=1}^n x_j^2 \right) \\ &= \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} p_{21} - p_{11} & \cdots & p_{\ell 1} - p_{11} & x_1 \\ p_{22} - p_{12} & \cdots & p_{\ell 2} - p_{12} & x_2 \\ \vdots & & \vdots & \vdots \\ \vdots & & \vdots & \vdots \\ p_{2n} - p_{1n} & \cdots & p_{\ell n} - p_{1n} & x_n \end{pmatrix}. \end{aligned}$$

Put

$$A = \begin{pmatrix} p_{21} - p_{11} & \cdots & p_{\ell 1} - p_{11} \\ p_{22} - p_{12} & \cdots & p_{\ell 2} - p_{12} \\ \vdots & & \vdots \\ \vdots & & \vdots \\ p_{2n} - p_{1n} & \cdots & p_{\ell n} - p_{1n} \end{pmatrix}.$$

Since ℓ -points $p_1, \dots, p_\ell$ are in general position, it is clearly seen that the rank of A is $\ell - 1$. Hence, by composing linear coordinate transformations if necessary, we may assume from the first that there exists an $(\ell - 1)$ by $(\ell - 1)$ regular matrix B such that

$$AB = \begin{pmatrix} 1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0} & & 1 \\ \alpha_{11} & \cdots & \alpha_{1,\ell-1} \\ \vdots & & \vdots \\ \alpha_{n-\ell+1,1} & \cdots & \alpha_{n-\ell+1,\ell-1} \end{pmatrix}.$$

Let $H_3 : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the diffeomorphism defined by

$$H_3(x_1, x_2, \dots, x_\ell) = \begin{pmatrix} x_1 & x_2 & \cdots & x_\ell \end{pmatrix} \left(\begin{array}{c|c} B & \begin{smallmatrix} 0 \\ \vdots \\ 0 \end{smallmatrix} \\ \hline 0 & \cdots & 0 & 1 \end{array} \right).$$

The composition of H_3 and $H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2$ is as follows.

$$\begin{aligned} & (H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2)(x_1, x_2, \dots, x_n) \\ &= \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} A & \begin{smallmatrix} x_1 \\ x_2 \\ \vdots \\ \vdots \\ x_n \end{smallmatrix} \end{pmatrix} \left(\begin{array}{c|c} B & \begin{smallmatrix} 0 \\ \vdots \\ 0 \end{smallmatrix} \\ \hline 0 & \cdots & 0 & 1 \end{array} \right) \\ &= \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} 1 & & \mathbf{0} & x_1 \\ & \ddots & & \vdots \\ \mathbf{0} & & 1 & x_{\ell-1} \\ \alpha_{11} & \cdots & \alpha_{1,\ell-1} & x_\ell \\ \vdots & & \vdots & \vdots \\ \alpha_{n-\ell+1,1} & \cdots & \alpha_{n-\ell+1,\ell-1} & x_n \end{pmatrix} \\ &= \left(x_1 + \sum_{j=1}^{n-\ell+1} \alpha_{j1} x_{\ell+j-1}, \dots, x_{\ell-1} + \sum_{j=1}^{n-\ell+1} \alpha_{j,\ell-1} x_{\ell+j-1}, \sum_{j=1}^n x_j^2 \right) \end{aligned}$$

Then, we have

$$\begin{aligned} & \left(x_1 + \sum_{j=1}^{n-\ell+1} \alpha_{j1} x_{\ell+j-1}, \dots, x_{\ell-1} + \sum_{j=1}^{n-\ell+1} \alpha_{j,\ell-1} x_{\ell+j-1}, x_{\ell}, \dots, x_n \right) \\ &= \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 0 & & & 0 \\ & & & 1 & & \\ \alpha_{11} & \cdots & \alpha_{1,\ell-1} & 1 & & \\ \vdots & & \vdots & 0 & \ddots & \\ \vdots & & \vdots & \vdots & & \ddots \\ \alpha_{n-\ell+1,1} & \cdots & \alpha_{n-\ell+1,\ell-1} & 0 & \cdots & 0 & 1 \end{pmatrix}. \end{aligned}$$

Put

$$C = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 0 & & & 0 \\ & & & 1 & & \\ \alpha_{11} & \cdots & \alpha_{1,\ell-1} & 1 & & \\ \vdots & & \vdots & 0 & \ddots & \\ \vdots & & \vdots & \vdots & & \ddots \\ \alpha_{n-\ell+1,1} & \cdots & \alpha_{n-\ell+1,\ell-1} & 0 & \cdots & 0 & 1 \end{pmatrix}.$$

It is easily seen that the inverse matrix of C is following:

$$C^{-1} = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 0 & & & 0 \\ & & & 1 & & \\ -\alpha_{11} & \cdots & -\alpha_{1,\ell-1} & 1 & & \\ \vdots & & \vdots & 0 & \ddots & \\ \vdots & & \vdots & \vdots & & \ddots \\ -\alpha_{n-\ell+1,1} & \cdots & -\alpha_{n-\ell+1,\ell-1} & 0 & \cdots & 0 & 1 \end{pmatrix}.$$

Let $H_4 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the diffeomorphism defined by

$$H_4(x) = xC^{-1},$$

where $x = (x_1, x_2, \dots, x_n)$. The composition of $H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2$ and H_4 is as follows:

$$\begin{aligned} & (H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2 \circ H_4)(x_1, x_2, \dots, x_n) \\ &= \left(x_1, x_2, \dots, x_{\ell-1}, \sum_{k=1}^{\ell-1} \left(x_k - \sum_{j=1}^{n-\ell+1} \alpha_{jk} x_{\ell+j-1} \right)^2 + \sum_{k=\ell}^n x_k^2 \right). \end{aligned}$$

Then, we have the following lemma.

Lemma 2.1. *There exist real numbers $r_j > 0$ ($1 \leq j \leq n$) and polynomials*

$$\varphi_j(x_1, \dots, x_{j-1}) = \sum_{k=1}^{j-1} a_{jk} x_k \quad (2 \leq j \leq n) \quad (a_{j1}, \dots, a_{j,j-1} \in \mathbb{R}) \text{ such that}$$

$$\sum_{k=1}^{\ell-1} \left(x_k - \sum_{j=1}^{n-\ell+1} \alpha_{jk} x_{\ell+j-1} \right)^2 + \sum_{k=\ell}^n x_k^2 = r_1 x_1^2 + \sum_{j=2}^n r_j (x_j + \varphi_j(x_1, \dots, x_{j-1}))^2.$$

The proof of Lemma 2.1 is given in Subsection 2.3. We continue to prove (I) of Proposition 2.1.

Let $H_5 : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the diffeomorphism defined by

$$\begin{aligned} & H_5(x_1, x_2, \dots, x_\ell) \\ &= \left(x_1, x_2, \dots, x_{\ell-1}, x_\ell - \left(r_1 x_1^2 + \sum_{j=2}^{\ell-1} r_j (x_j + \varphi_j(x_1, \dots, x_{j-1}))^2 \right) \right). \end{aligned}$$

The composition of H_5 and $H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2 \circ H_4$ is as follows:

$$\begin{aligned} & (H_5 \circ H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2 \circ H_4)(x_1, x_2, \dots, x_n) \\ &= \left(x_1, x_2, \dots, x_{\ell-1}, \sum_{j=\ell}^n r_j (x_j + \varphi_j(x_1, \dots, x_{j-1}))^2 \right). \end{aligned}$$

Let $H_6 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the diffeomorphism defined by

$$\begin{aligned} & H_6(x_1, x_2, \dots, x_n) \\ &= \left(x_1, \dots, x_{\ell-1}, \frac{x_\ell}{\sqrt{r_\ell}} - \varphi_\ell(x_1, \dots, x_{\ell-1}), \dots, \frac{x_n}{\sqrt{r_n}} - \varphi_n(x_1, \dots, x_{n-1}) \right). \end{aligned}$$

It follows that

$$\begin{aligned} & (H_5 \circ H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2 \circ H_4 \circ H_6)(x_1, x_2, \dots, x_n) \\ &= (x_1, x_2, \dots, x_{\ell-1}, x_\ell^2 + \dots + x_n^2). \end{aligned}$$

□

2.2. Proof of (II). Similarly as proof of (I) of Proposition 2.1, the composition $H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2$ is as follows:

$$\begin{aligned} & (H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2)(x_1, x_2, \dots, x_n) \\ &= \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} p_{21} - p_{11} & \cdots & p_{\ell 1} - p_{11} & x_1 \\ p_{22} - p_{12} & \cdots & p_{\ell 2} - p_{12} & x_2 \\ \vdots & & \vdots & \vdots \\ \vdots & & \vdots & \vdots \\ p_{2n} - p_{1n} & \cdots & p_{\ell n} - p_{1n} & x_n \end{pmatrix}, \end{aligned}$$

and put

$$A = \begin{pmatrix} p_{21} - p_{11} & \cdots & p_{\ell 1} - p_{11} \\ p_{22} - p_{12} & \cdots & p_{\ell 2} - p_{12} \\ \vdots & & \vdots \\ \vdots & & \vdots \\ p_{2n} - p_{1n} & \cdots & p_{\ell n} - p_{1n} \end{pmatrix}.$$

Since $(n+1)$ -points $p_1, \dots, p_{n+1}$ are in general position, it is clearly seen that the rank of A is n . Hence, there exists an $(\ell-1)$ by $(\ell-1)$ regular matrix B such that

$$AB = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ & \ddots & \vdots & & \vdots \\ 0 & & 1 & 0 & \cdots & 0 \end{pmatrix}.$$

Similarly as the proof of (I) of Proposition 2.1, the composition $H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2$ is as follows:

$$\begin{aligned} & (H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2)(x_1, x_2, \dots, x_n) \\ &= (x_1 \ x_2 \ \cdots \ x_n) \begin{pmatrix} A & \begin{array}{c} x_1 \\ x_2 \\ \vdots \\ x_n \end{array} \end{pmatrix} \begin{pmatrix} B & \begin{array}{c} 0 \\ \vdots \\ 0 \end{array} \\ \hline 0 & \cdots & 0 & 1 \end{pmatrix} \\ &= (x_1 \ x_2 \ \cdots \ x_n) \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & x_1 \\ & \ddots & \vdots & & \vdots & \vdots \\ 0 & & 1 & 0 & \cdots & 0 & x_n \end{pmatrix} \\ &= \left(x_1, \dots, x_n, 0, \dots, 0, \sum_{j=1}^n x_j^2 \right). \end{aligned}$$

Let $\tilde{H}_4 : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the diffeomorphism defined by

$$\tilde{H}_4(x_1, x_2, \dots, x_\ell) = \left(x_1, \dots, x_n, \dots, x_{\ell-1}, x_\ell - \sum_{j=1}^n x_j^2 \right).$$

It follows that

$$\begin{aligned} & (\tilde{H}_4 \circ H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)} \circ H_2)(x_1, x_2, \dots, x_n) \\ &= (x_1, x_2, \dots, x_n, 0, \dots, 0). \end{aligned}$$

□

2.3. Proof of Lemma 2.1. Put

$$(1) \quad f(x_1, x_2, \dots, x_n) = \sum_{k=1}^{\ell-1} \left(x_k - \sum_{j=1}^{n-\ell+1} \alpha_{jk} x_{\ell+j-1} \right)^2 + \sum_{k=\ell}^n x_k^2.$$

Since $f(x_1, x_2, \dots, x_n)$ is a quadratic form of $x_1, x_2, \dots, x_n$, it is not hard to see that there exist real numbers r_j ($1 \leq j \leq n$) and polynomials

$$\varphi_j(x_1, \dots, x_{j-1}) = \sum_{k=1}^{j-1} a_{jk} x_k \quad (2 \leq j \leq n) \quad (a_{j1}, \dots, a_{j,j-1} \in \mathbb{R})$$

such that

$$(2) \quad f(x_1, x_2, \dots, x_n) = r_1 x_1^2 + \sum_{j=2}^n r_j (x_j + \varphi_j(x_1, \dots, x_{j-1}))^2.$$

Next, we prove that $r_j > 0$ for any j ($\ell \leq j \leq n$). Suppose that there exists a natural number j ($\ell \leq j \leq n$) such that $r_j \leq 0$. Put $x_1 = \cdots = x_{j-1} = 0$, and we get $\varphi_2(0) = \cdots = \varphi_j(0, \dots, 0) = 0$. Then, put

$$\begin{aligned} x_j &= \varepsilon_j (\neq 0) \\ x_{j+1} &= -\varphi_{j+1}(0, \dots, 0, \varepsilon_j) = \varepsilon_{j+1} \\ x_{j+2} &= -\varphi_{j+2}(0, \dots, 0, \varepsilon_j, \varepsilon_{j+1}) = \varepsilon_{j+2} \\ &\vdots \\ x_n &= -\varphi_n(0, \dots, 0, \varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_{n-1}) = \varepsilon_n. \end{aligned}$$

By (2), we get $f(0, \dots, 0, \varepsilon_j, \dots, \varepsilon_n) = r_j \varepsilon_j^2$. Since $r_j \leq 0$ and $\varepsilon_j^2 > 0$, we obtain $f(0, \dots, 0, \varepsilon_j, \dots, \varepsilon_n) \leq 0$.

On the other hand, by (1), it is clearly seen that $f(0, \dots, 0, \varepsilon_j, \dots, \varepsilon_n) \geq \varepsilon_j^2 > 0$. This is a contradiction. Hence, it follows that $r_j > 0$ for any j ($\ell \leq j \leq n$). $\square$

3. PROOFS OF THEOREMS 1.1 AND 1.2

Frist, we prepare the following lemma.

Lemma 3.1 ([2, 3]). *Let M be a closed manifold and let $i : M \rightarrow \mathbb{R}^\ell$ be an immersion with normal crossing. Then, there exists a subset $\Sigma \subset \mathcal{L}(\mathbb{R}^\ell, \mathbb{R})$ of Lebesgue measure 0 such that for any $\pi \in \mathcal{L}(\mathbb{R}^\ell, \mathbb{R}) - \Sigma$, the composition $\pi \circ i : M \rightarrow \mathbb{R}$ is a Morse function, where $\mathcal{L}(\mathbb{R}^\ell, \mathbb{R})$ stands for the space of linear mappings $\mathbb{R}^\ell \rightarrow \mathbb{R}$.*

3.1. Proof of Theorem 1.1. Since M is a closed manifold, and $i : M \rightarrow \mathbb{R}^\ell$ is an embedding, by Lemma 3.1 there exists a linear mapping $\pi : \mathbb{R}^\ell \rightarrow \mathbb{R}$:

$$\pi(x) = \sum_{j=1}^{\ell} u_j x_j \quad (u = (u_1, \dots, u_\ell), x = (x_1, \dots, x_\ell))$$

such that $\pi \circ i : M \rightarrow \mathbb{R}$ is a Morse function. By transforming basis of the linear space $\mathbb{R}^\ell$ if necessary, we may assume from the first that $u = (0, \dots, 0, 1)$. Note that $\pi(x) = x_\ell$, $\pi \circ i = i_\ell$ ($i = (i_1, \dots, i_\ell)$). Since M is compact and $\pi \circ i$ is continuous, $\pi \circ i$ has the minimal value. Let k_0 be the minimal value of $\pi \circ i$. It is clearly seen that k_0 is a critical value. Since $\pi \circ i$ is a Morse function, the set $(\pi \circ i)^{-1}(k_0)$ consists of only one point. Let q be the point such that $(\pi \circ i)(q) = k_0$. Let $(U, \varphi, (t_1, \dots, t_m))$ be a coordinate neighborhood of q .

Let $(Ji)_q$ be the Jacobian matrix of the mapping i at q . Note that

$$\frac{\partial(i_\ell \circ \varphi^{-1})}{\partial t_j}(\varphi(q)) = 0$$

for any j ($1 \leq j \leq m$). Thus, we have

$$(0, \dots, 0, 1)(Ji)_q = (0, \dots, 0).$$

By transforming basis of the linear space $\mathbb{R}^\ell$ again if necessary, we may assume from the first that the following matrix is invertible:

$$(E_m \mid O)(Ji)_q.$$

Here, E_m stands for the m by m unit matrix and O stands for the m by $(\ell - m)$ zero matrix.

Put $A = \mathbb{R}^m \times \{0\} \subset \mathbb{R}^\ell$. Consider the mapping $\psi : \varphi(U) \rightarrow A$ defined by

$$\psi = (i_1 \circ \varphi^{-1}, \dots, i_m \circ \varphi^{-1}).$$

It is clearly seen that $(J\psi)_{\varphi(q)}$ is invertible. By the inverse function theorem, there exist an open neighborhood V of $\varphi(q)$ and an open neighborhood $\tilde{V}$ of $\psi(\varphi(q))$ such that $\psi : V \rightarrow \tilde{V}$ is a diffeomorphism. We may suppose that V and $\tilde{V}$ are connected and $V \subset \varphi(U)$. Note that $i(\varphi^{-1}(V))$ can be expressed as follows:

$$i(\varphi^{-1}(V)) = \{(x, \xi_{m+1}(x), \dots, \xi_\ell(x)) \in \mathbb{R}^\ell \mid x \in \tilde{V}\},$$

where $x = (x_1, \dots, x_m)$ and $\xi_j = i_j \circ \varphi^{-1} \circ \psi^{-1}$ for any j ($m+1 \leq j \leq \ell$). Put

$$k_1 = \sup_{q' \in \varphi^{-1}(V)} i_\ell(q').$$

It is not hard to see that $k_1 > k_0$, and that $i_\ell^{-1}(r)$ is a non-empty set for any $r \in [k_0, k_1]$. Since $M - \varphi^{-1}(V)$ is compact and i_ℓ is continuous, there exists a minimal value in $i_\ell(M - \varphi^{-1}(V))$. Let k_2 be the minimal value of $i_\ell(M - \varphi^{-1}(V))$. Then, it is clearly seen that $k_2 > k_0$.

Put

$$k_3 = \min\left\{k_0 + 1, k_0 + \frac{k_2 - k_0}{3}, k_1\right\}.$$

Then, we have the following lemma.

Lemma 3.2. *There exists a real number $a \in (k_0, k_3)$ such that there exist $(m+1)$ -points in general position in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$.*

The proof of Lemma 3.2 is given in Subsection 3.3. We continue to prove Theorem 1.1. Let $p_1, \dots, p_{m+1}$ be $(m+1)$ -points in general position in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Then, we take $\ell - (m+1)$ -points $p_{m+2}, \dots, p_\ell \in \mathbb{R}^{\ell-1} \times \{a\}$ such that ℓ -points $p_1, \dots, p_\ell$ are in general position.

Note that $p_{1\ell} = \dots = p_{\ell\ell} = a$. Similarly as the proof of Proposition 2.1, the composition $H_1 \circ D_{(p_1, \dots, p_\ell)}$ can be expressed as follows:

$$\begin{aligned} & (H_1 \circ D_{(p_1, \dots, p_\ell)})(x_1, x_2, \dots, x_\ell) \\ &= \begin{pmatrix} x_1 - p_{11} \\ \vdots \\ \vdots \\ x_\ell - a \end{pmatrix}^t \begin{pmatrix} p_{21} - p_{11} & \cdots & p_{\ell 1} - p_{11} & x_1 - p_{11} \\ p_{22} - p_{12} & \cdots & p_{\ell 2} - p_{12} & x_2 - p_{12} \\ \vdots & & \vdots & \vdots \\ p_{2, \ell-1} - p_{1, \ell-1} & \cdots & p_{\ell, \ell-1} - p_{1, \ell-1} & x_{\ell-1} - p_{1, \ell-1} \\ 0 & \cdots & 0 & x_\ell - a \end{pmatrix}, \end{aligned}$$

where ${}^t A$ means a transposed matrix of A . The composition of H_3 and $H_1 \circ D_{(p_1, \dots, p_\ell)}$ is as follows:

$$\begin{aligned} & (H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)})(x_1, x_2, \dots, x_\ell) \\ &= \left(x_1 - p_{11}, \dots, x_{\ell-1} - p_{1, \ell-1}, \sum_{j=1}^{\ell-1} (x_j - p_{1j})^2 + (x_\ell - a)^2 \right). \end{aligned}$$

Let $H'_4 : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the diffeomorphism defined by

$$H'_4(x_1, x_2, \dots, x_\ell) = \left(x_1 + p_{11}, \dots, x_{\ell-1} + p_{1, \ell-1}, x_\ell - \sum_{j=1}^{\ell-1} x_j^2 + a \right).$$

Then, we have the following:

$$\begin{aligned} & (H'_4 \circ H_3 \circ H_1 \circ D_{(p_1, \dots, p_\ell)})(x_1, x_2, \dots, x_\ell) \\ &= (x_1, x_2, \dots, x_{\ell-1}, (x_\ell - a)^2 + a). \end{aligned}$$

For simplicity, put $H = H'_4 \circ H_3 \circ H_1$. Since H is a diffeomorphism, in order to prove that $D_{(p_1, \dots, p_\ell)} \circ i$ is an embedding, it is sufficient to show that $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is an embedding. The mapping $H \circ D_{(p_1, \dots, p_\ell)}$ is a definite fold mapping whose singular set is the hyperplane $\mathbb{R}^{\ell-1} \times \{a\}$. Hence, any singular point of $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ must be contained in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Since the intersection $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$ is contained in $i(\varphi^{-1}(V))$, the following equality holds for any point in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$:

$$\begin{aligned} & (H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}(x, x_{m+1}, \dots, x_\ell) \\ &= (H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}(x, \xi_{m+1}(x), \dots, \xi_\ell(x)) \\ &= (x, \xi_{m+1}(x), \dots, \xi_{\ell-1}(x), (\xi_\ell(x) - a)^2 + a), \end{aligned}$$

where $x = (x_1, \dots, x_m)$. Note that the rank of the Jacobian matrix of $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ at any point of $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$ is always m , which implies that $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is non-singular.

Then, we have the following lemma.

Lemma 3.3. $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is injective.

The proof of Lemma 3.3 is given in Subsection 3.4. We continue to prove Theorem 1.1. Since $i(M)$ is compact, $\mathbb{R}^\ell$ is a Hausdorff space, and $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is continuous and injective, we see that $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is homeomorphic to the image. Since i is an embedding, $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is an embedding. $\square$

3.2. Proof of Theorem 1.2. Since M is a closed manifold, and $i : M \rightarrow \mathbb{R}^\ell$ is an immersion with normal crossing, by Lemma 3.1, there exists a linear mapping $\pi : \mathbb{R}^\ell \rightarrow \mathbb{R}$:

$$\pi(x) = \sum_{j=1}^{\ell} u_j x_j \quad (u = (u_1, \dots, u_\ell), x = (x_1, \dots, x_\ell))$$

such that $\pi \circ i : M \rightarrow \mathbb{R}$ is a Morse function. By transforming basis of the linear space $\mathbb{R}^\ell$ if necessary, we may assume from the first that $u = (0, \dots, 0, 1)$. Note that $\pi(x) = x_\ell$, $\pi \circ i = i_\ell$ ($i = (i_1, \dots, i_\ell)$). Since M is compact and $\pi \circ i$ is continuous, $\pi \circ i$ has the minimal value. Let k_0 be the minimal value of $\pi \circ i$. It is clearly seen that k_0 is a critical value. Since $\pi \circ i$ is a Morse function, the set $(\pi \circ i)^{-1}(k_0)$ consists of only one point. Let q be the point such that $(\pi \circ i)(q) = k_0$. Let $(U, \varphi, (t_1, \dots, t_m))$ be a coordinate neighborhood of q . By shrinking U sufficiently small if necessary, we may assume that $i|_U : U \rightarrow \mathbb{R}^\ell$ is an embedding. Let $(Ji)_q$ be the Jacobian matrix of the mapping i at q . Note that

$$\frac{\partial(i_\ell \circ \varphi^{-1})}{\partial t_j}(\varphi(q)) = 0$$

for any j ($1 \leq j \leq m$). Thus, we have

$$(0, \dots, 0, 1)(Ji)_q = (0, \dots, 0).$$

By transforming basis of the linear space $\mathbb{R}^\ell$ again if necessary, we may assume from the first that the following matrix is invertible:

$$(E_m \mid O)(Ji)_q.$$

Here, E_m stands for the m by m unit matrix and O stands for the m by $(\ell - m)$ zero matrix.

Put $A = \mathbb{R}^m \times \{0\} \subset \mathbb{R}^\ell$. Consider the mapping $\psi : \varphi(U) \rightarrow A$ defined by

$$\psi = (i_1 \circ \varphi^{-1}, \dots, i_m \circ \varphi^{-1}).$$

It is clearly seen that $(J\psi)_{\varphi(q)}$ is invertible. By the inverse function theorem, there exist an open neighborhood V of $\varphi(q)$ and an open neighborhood $\tilde{V}$ of $\psi(\varphi(q))$ such that $\psi : V \rightarrow \tilde{V}$ is a diffeomorphism. We may suppose that V and $\tilde{V}$ are connected and $V \subset \varphi(U)$. Note that $i(\varphi^{-1}(V))$ can be expressed as follows:

$$i(\varphi^{-1}(V)) = \{(x, \xi_{m+1}(x), \dots, \xi_\ell(x)) \in \mathbb{R}^\ell \mid x \in \tilde{V}\},$$

where $x = (x_1, \dots, x_m)$ and $\xi_j = i_j \circ \varphi^{-1} \circ \psi^{-1}$ for any j ($m+1 \leq j \leq \ell$). Put

$$k_1 = \sup_{q' \in \varphi^{-1}(V)} i_\ell(q').$$

It is not hard to see that $k_1 > k_0$, and that $i_\ell^{-1}(r)$ is a non-empty set for any $r \in [k_0, k_1]$. Since $M - \varphi^{-1}(V)$ is compact and i_ℓ is continuous, there exist a minimal value in $i_\ell(M - \varphi^{-1}(V))$. Let k_2 be the minimal value of $i_\ell(M - \varphi^{-1}(V))$. Then, it is clearly seen that $k_2 > k_0$.

Put

$$k_3 = \min\left\{k_0 + 1, k_0 + \frac{k_2 - k_0}{3}, k_1\right\}.$$

By Lemma 3.2, there exists a real number $a \in (k_0, k_3)$ such that there exist $(m+1)$ -points in general position in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Let $p_1, \dots, p_{m+1}$ be $(m+1)$ -points in general position in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Then, we take $\ell - (m+1)$ -points $p_{m+2}, \dots, p_\ell \in \mathbb{R}^{\ell-1} \times \{a\}$ such that ℓ -points $p_1, \dots, p_\ell$ are in general position.

Then, by the diffeomorphism $H : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ defined in Subsection 3.1, we have the following:

$$(H \circ D_{(p_1, \dots, p_\ell)})(x_1, x_2, \dots, x_\ell) = (x_1, x_2, \dots, x_{\ell-1}, (x_\ell - a)^2 + a).$$

Since H is a diffeomorphism, in order to prove that $D_{(p_1, \dots, p_\ell)} \circ i$ is an immersion with normal crossing, it is sufficient to show that $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is an immersion with normal crossing. Then, similarly as the proof of Theorem 1.1, it is seen that $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is non-singular, which implies that $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is an immersion.

Finally, we prove that $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is a mapping with normal crossing. Since by Lemma 3.3 $(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)}$ is injective, for any $y \in (H \circ D_{(p_1, \dots, p_\ell)} \circ i)(M)$, the set $((H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)})^{-1}(y)$ has only one point. Let x be the only one point.

Suppose that $x \in i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Since $i^{-1}(x) \subset \varphi^{-1}(V)$ and $i|_{\varphi^{-1}(V)} : \varphi^{-1}(V) \rightarrow \mathbb{R}^\ell$ is an embedding, the set $i^{-1}(x)$ has only one point. Hence, it is clearly seen that $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is a mapping with normal crossing at the point.

Suppose that $x \in i(M) - (\mathbb{R}^{\ell-1} \times \{a\})$. Since i is a mapping with normal crossing, the set $(H \circ D_{(p_1, \dots, p_\ell)} \circ i)^{-1}(y)$ can be expressed as follows:

$$(H \circ D_{(p_1, \dots, p_\ell)} \circ i)^{-1}(y) = i^{-1}(x) = \{q_1, \dots, q_n\}.$$

Since $d(H \circ D_{(p_1, \dots, p_\ell)})_x : T_x \mathbb{R}^\ell \rightarrow T_y \mathbb{R}^\ell$ is a bijective linear mapping, for any subset $\{\lambda_1, \lambda_2, \dots, \lambda_s\} \subset \{1, 2, \dots, n\}$ ($s \leq n$), the following equality holds.

$$\begin{aligned} & \text{codim} \left(\bigcap_{j=1}^s d(H \circ D_{(p_1, \dots, p_\ell)} \circ i)_{q_{\lambda_j}}(T_{q_{\lambda_j}} M) \right) \\ &= \ell - \dim \bigcap_{j=1}^s d(H \circ D_{(p_1, \dots, p_\ell)} \circ i)_{q_{\lambda_j}}(T_{q_{\lambda_j}} M) \\ &= \ell - \dim d(H \circ D_{(p_1, \dots, p_\ell)})_x \left(\bigcap_{j=1}^s di_{q_{\lambda_j}}(T_{q_{\lambda_j}} M) \right) \\ &= \ell - \dim \left(\bigcap_{j=1}^s di_{q_{\lambda_j}}(T_{q_{\lambda_j}} M) \right) \\ &= \sum_{j=1}^s (\ell - \dim di_{q_{\lambda_j}}(T_{q_{\lambda_j}} M)) \\ &= \sum_{j=1}^s (\ell - \dim d(H \circ D_{(p_1, \dots, p_\ell)} \circ i)_{q_{\lambda_j}}(T_{q_{\lambda_j}} M)) \\ &= \sum_{j=1}^s \text{codim} d(H \circ D_{(p_1, \dots, p_\ell)} \circ i)_{q_{\lambda_j}}(T_{q_{\lambda_j}} M) \end{aligned}$$

Hence, $H \circ D_{(p_1, \dots, p_\ell)} \circ i$ is a mapping with normal crossing. $\square$

3.3. Proof of Lemma 3.2. First, suppose that $m \geq 2$. By Sard's theorem, there exists a real number $a \in (k_0, k_3)$ such that a is a regular value of the mapping $i_\ell : M \rightarrow \mathbb{R}$. Since a real number a is also an element of (k_0, k_1) , we get $i_\ell^{-1}(a) \neq \emptyset$. Hence, $i_\ell^{-1}(a)$ is an $(m-1)$ -dimensional closed manifold. Note that $i(i_\ell^{-1}(a)) = i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$. Let n_0 be the maximal value of the number of points which are in general position in $i(i_\ell^{-1}(a))$. It is clearly seen that $n_0 \geq 2$. Let $p_1, \dots, p_{n_0}$ be n_0 -points in general position in $i(i_\ell^{-1}(a))$. Put

$$V = \left\{ \sum_{j=2}^{n_0} \alpha_j \overrightarrow{p_1 p_j} \mid \alpha_j \in \mathbb{R} \right\}.$$

It is not hard to see that $i(i_\ell^{-1}(a)) \subset V$ and $\dim V = n_0 - 1$. Since $i|_{i_\ell^{-1}(a)} : i_\ell^{-1}(a) \rightarrow V$ is an immersion, and $i|_{i_\ell^{-1}(a)}$ is closed, it follows that $m-1 < n_0-1$.

Next, suppose that $m = 1$. Since $\tilde{V}$ is a bounded connected open set of $\mathbb{R}$, we can put $\tilde{V} = (b_1, b_2)$. Then, the set $i(\varphi^{-1}(V))$ can be expressed as follows:

$$i(\varphi^{-1}(V)) = \{(x, \xi_2(x), \dots, \xi_\ell(x)) \in \mathbb{R}^\ell \mid x \in (b_1, b_2)\}.$$

Note that $b_1 < i_1(q) < b_2$, and define two closed intervals as follows:

$$\begin{aligned} B_1 &= \left[b_1 + \frac{i_1(q) - b_1}{2}, i_1(q) \right], \\ B_2 &= \left[i_1(q), i_1(q) + \frac{b_2 - i_1(q)}{2} \right]. \end{aligned}$$

It is not hard to see that

$$\begin{aligned} \xi_\ell \left(b_1 + \frac{i_1(q) - b_1}{2} \right) &> \xi_\ell(i_1(q)) = k_0, \\ \xi_\ell \left(i_1(q) + \frac{b_2 - i_1(q)}{2} \right) &> \xi_\ell(i_1(q)) = k_0. \end{aligned}$$

Then, put

$$a = \min \left\{ \xi_\ell \left(b_1 + \frac{i_1(q) - b_1}{2} \right), \xi_\ell \left(i_1(q) + \frac{b_2 - i_1(q)}{2} \right), \frac{k_0 + k_3}{2} \right\}.$$

It is clearly seen that $a \in (k_0, k_3)$. Since B_1 and B_2 are compact and ξ_ℓ is continuous, by intermediate value theorem, there exists a real number $c_1 \in B_1$ such that $\xi_\ell(c_1) = a$ and there exists a real number $c_2 \in B_2$ such that $\xi_\ell(c_2) = a$. Since it is clearly seen that $c_1 \neq c_2$, there exist two points $(c_1, \xi_2(c_1), \dots, \xi_\ell(c_1))$, $(c_2, \xi_2(c_2), \dots, \xi_\ell(c_2))$ in $i(M) \cap (\mathbb{R}^{\ell-1} \times \{a\})$ at least. $\square$

3.4. Proof of Lemma 3.3. For a real number r , put

$$\begin{aligned} A_{<r} &= i(M) \cap (\mathbb{R}^{\ell-1} \times \{x \in \mathbb{R} \mid x < r\}), \\ A_{\geq r} &= i(M) \cap (\mathbb{R}^{\ell-1} \times \{x \in \mathbb{R} \mid x \geq r\}). \end{aligned}$$

Since $H \circ D_{(p_1, \dots, p_\ell)}$ is a definite fold mapping whose singular set is the hyperplane $\mathbb{R}^{\ell-1} \times \{a\}$, the following can be obtained easily:

$$\begin{aligned} &(H \circ D_{(p_1, \dots, p_\ell)})|_{i(M)} \text{ is injective} \\ \Leftrightarrow &(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{\geq a} = \emptyset. \end{aligned}$$

In order to prove $(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{\geq a} = \emptyset$, it is sufficient to show the following (1) and (2).

$$\begin{aligned} (1) \quad &(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{\geq k_2} = \emptyset \\ (2) \quad &(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{<k_2} = \emptyset \end{aligned}$$

First, we prove (1). Suppose that $(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{\geq k_2} \neq \emptyset$. Then, there exists a real number $\tilde{a} \in [k_0, a)$ such that $(\tilde{a} - a)^2 + a \geq k_2$. It is clearly seen that $(k_0 - a)^2 + a \geq (\tilde{a} - a)^2 + a$. Hence, we get $(k_0 - a)^2 + a \geq k_2$. On the other hand, since $k_0 < a < k_0 + 1$, we obtain $0 < a - k_0 < 1$. Thus, we have $(k_0 - a)^2 + a < 2(a - k_0) + a$. Since $a < k_0 + (k_2 - k_0)/3$, we get $2(a - k_0) + a < k_2$. Therefore, we obtain $(k_0 - a)^2 + a < k_2$. However, this is inconsistent with $(k_0 - a)^2 + a \geq k_2$.

Next, we prove (2). Suppose that $(H \circ D_{(p_1, \dots, p_\ell)})(A_{<a}) \cap A_{<k_2} \neq \emptyset$. Note that $A_{<k_2} \subset i(\varphi^{-1}(V))$, and it is clearly seen that there exist two points

$$(x, \xi_{m+1}(x), \dots, \xi_\ell(x)) \in A_{<a}, \quad (x', \xi_{m+1}(x'), \dots, \xi_\ell(x')) \in A_{<k_2}$$

such that

$$(H \circ D_{(p_1, \dots, p_\ell)})(x, \xi_{m+1}(x), \dots, \xi_\ell(x)) = (x', \xi_{m+1}(x'), \dots, \xi_\ell(x')),$$

where $x = (x_1, \dots, x_m)$ and $x' = (x'_1, \dots, x'_m)$. Thus, since $x = x'$ and $\xi_\ell(x') = (\xi_\ell(x) - a)^2 + a$, we get $\xi_\ell(x) \geq a$. However, this contradicts to $\xi_\ell(x) < a$. $\square$

4. PROOF OF PROPOSITION 1.1

4.1. **Preparations.** First, we prepare the following three lemmas.

Lemma 4.1 ([4], p.354). *Let $A \subset \mathbb{R}^2$ be a closed convex set, and let x be a point in ∂A , where ∂A means a boundary of A . Then, there exist the following sets*

$$\begin{aligned} L &= \{(x_1, x_2) \in \mathbb{R}^2 \mid \alpha x_1 + \beta x_2 + \gamma = 0\}, \\ L_{\leq} &= \{(x_1, x_2) \in \mathbb{R}^2 \mid \alpha x_1 + \beta x_2 + \gamma \leq 0\} \end{aligned}$$

such that $x \in L$, $A \subset L_{\leq}$ ($\alpha \neq 0 \vee \beta \neq 0$).

Lemma 4.2 ([4], p.339). *Let A be a subset of $\mathbb{R}^2$. Then, for any point $x = (x_1, x_2) \in \text{conv}(A)$, there exist three points $q_1 = (q_{11}, q_{12})$, $q_2 = (q_{21}, q_{22})$, $q_3 = (q_{31}, q_{32}) \in A$ and real numbers t_1, t_2, t_3 such that*

$$\begin{aligned} x_1 &= t_1 q_{11} + t_2 q_{21} + t_3 q_{31} \\ x_2 &= t_1 q_{12} + t_2 q_{22} + t_3 q_{32} \\ (t_1 \geq 0, t_2 \geq 0, t_3 \geq 0, t_1 + t_2 + t_3 &= 1), \end{aligned}$$

where $\text{conv}(A)$ means a convex hull of A .

Lemma 4.3 ([4], p.343). *Let A be a subset of $\mathbb{R}^2$ homeomorphic to S^1 . Then, $\partial \text{conv}(A)$ is homeomorphic to S^1 .*

Now, we define the following mappings which are important in the proof of Proposition 1.1.

Definition 4.1. Let $\pi_{\theta} : i(S^1) \rightarrow \mathbb{R}$ be the mapping defined by

$$\pi_{\theta}(x_1, x_2) = (\cos \theta)x_1 + (\sin \theta)x_2.$$

Since $i(S^1)$ is compact and π_{θ} is continuous, π_{θ} has the maximal value. Let k_{θ} be the maximal value of π_{θ} . Then, it is clearly seen that we obtain (I) or (II) as follows.

(I) For any $\theta \in [0, 2\pi)$, the set $\pi_{\theta}^{-1}(k_{\theta})$ has only one point.

(II) There exists $\theta \in [0, 2\pi)$ such that the set $\pi_{\theta}^{-1}(k_{\theta})$ has two points at least.

Then, we have the following lemma.

Lemma 4.4. *In the circumstance of (I), $i(S^1) = \partial \text{conv}(i(S^1))$.*

Proof of Lemma 4.4.

First, we prove that $\partial \text{conv}(i(S^1)) \subset i(S^1)$. Suppose that there exists $a \in \partial \text{conv}(i(S^1))$ such that $a \notin i(S^1)$ ($a = (a_1, a_2)$). Then, by Lemma 4.1, there exist the following sets:

$$\begin{aligned} L &= \{(x_1, x_2) \in \mathbb{R}^2 \mid \alpha x_1 + \beta x_2 + \gamma = 0\}, \\ L_{\leq} &= \{(x_1, x_2) \in \mathbb{R}^2 \mid \alpha x_1 + \beta x_2 + \gamma \leq 0\} \end{aligned}$$

such that $a \in L$, $\text{conv}(i(S^1)) \subset L_{\leq}$. By transforming the origin and basis of the linear space $\mathbb{R}^2$ if necessary, we may assume from the first that

$$\begin{aligned} a &= (0, 0), \\ L &= \mathbb{R} \times \{0\}, \\ L_{\leq} &= \mathbb{R} \times \{x \in \mathbb{R} \mid x \leq 0\}. \end{aligned}$$

Note that $(a_1, a_2) = (0, 0)$, and by Lemma 4.2 there exist three points $q_1 = (q_{11}, q_{12})$, $q_2 = (q_{21}, q_{22})$, $q_3 = (q_{31}, q_{32}) \in i(S^1)$ and real numbers t_1, t_2, t_3 such that

$$(3) \quad t_1 q_{11} + t_2 q_{21} + t_3 q_{31} = 0$$

$$(4) \quad t_1 q_{12} + t_2 q_{22} + t_3 q_{32} = 0$$

$$(t_1 \geq 0, t_2 \geq 0, t_3 \geq 0, t_1 + t_2 + t_3 = 1).$$

Note that $q_{12} \leq 0$, $q_{22} \leq 0$, $q_{32} \leq 0$, and it is not hard to see that there exists j ($1 \leq j \leq 3$) such that $q_{j2} = 0$. Then, we may assume that $q_{12} = 0$.

Thus, we see that the maximal value of the mapping $\pi_{\pi/2}$ is 0. Since the set $\pi_{\pi/2}^{-1}(0)$ has only one point, it is clearly seen that $q_{22} < 0$, $q_{32} < 0$. Hence, by (2), it is necessary to be $t_2 = t_3 = 0$. Then, since we have $t_1 = 1$, we obtain $a = q_1$ by (1) and (2). This is inconsistent with $a \notin i(S^1)$.

Next, we prove that $i(S^1) \subset \partial \text{conv}(i(S^1))$. Suppose that there exists $a \in i(S^1)$ such that $a \notin \partial \text{conv}(i(S^1))$. Since $\partial \text{conv}(i(S^1)) \subset i(S^1)$, we obtain $\partial \text{conv}(i(S^1)) \subset i(S^1) - \{a\}$. Then, we see that it means that $\partial \text{conv}(i(S^1))$ is not homeomorphic to S^1 . This is inconsistent with Lemma 4.3 $\square$

4.2. Proof. Since $i(S^1)$ is compact, $\mathbb{R}^2$ is a Hausdorff space, $d_{(p_1, p_2)}|_{i(S^1)}$ is continuous, and i is homeomorphic to the image, in order to prove that $d_{(p_1, p_2)} \circ i$ is homeomorphic to the image, it is sufficient to show that $d_{(p_1, p_2)}|_{i(S^1)}$ is injective.

First, in the circumstance of (I), we prove that $d_{(p_1, p_2)}|_{i(S^1)}$ is injective. Let $d : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be the two-dimensional Euclidean distance defined in Section 1. Since the set $i(S^1) \times i(S^1)$ is compact and d is continuous, $d|_{i(S^1) \times i(S^1)}$ has a maximal value. Let k be the maximal value of $d|_{i(S^1) \times i(S^1)}$. Let $(a, a') \in i(S^1) \times i(S^1)$ be an element of the set $(d|_{i(S^1) \times i(S^1)})^{-1}(k)$. By transforming the origin and basis of the linear space $\mathbb{R}^2$ if necessary, we may assume from the first that $a = (0, 0)$, $a' = (1, 0)$. Then, note that $k = 1$.


Put $\theta_0 = \pi/2$, and define $\pi_{\theta_0} : i(S^1) \rightarrow \mathbb{R}$; thus

$$\pi_{\theta_0}(x_1, x_2) = x_2.$$

Since $i(S^1)$ is compact and π_{θ_0} is continuous, π_{θ_0} has a maximal value and a minimal value. Let k_{θ_0} be the maximal value of the mapping π_{θ_0} , and let $\tilde{k}_{\theta_0}$ be the minimal value of the mapping π_{θ_0} . Since there exist two points $a = (0, 0)$, $a' = (1, 0)$ in $i(S^1)$, it is clearly seen that $\tilde{k}_{\theta_0} \leq 0 \leq k_{\theta_0}$. Then, the set $\pi_{\theta_0}^{-1}(k_{\theta_0})$ has only one point. Since $-\tilde{k}_{\theta_0}$ is also the maximal value of $\pi_{3\pi/2}$ and $\pi_{\theta_0}^{-1}(\tilde{k}_{\theta_0}) = \pi_{3\pi/2}^{-1}(-\tilde{k}_{\theta_0})$, the set $\pi_{\theta_0}^{-1}(\tilde{k}_{\theta_0})$ also has only one point. Hence, it is not hard to see that $\tilde{k}_{\theta_0} < 0 < k_{\theta_0}$. By transforming basis of the linear space $\mathbb{R}^2$ again if necessary, we may assume from the first that $|\tilde{k}_{\theta_0}| \leq |k_{\theta_0}|$.

We shall prove that the set $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3)$ has two points at least. Note that $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3) = i(S^1) \cap (\mathbb{R} \times \{2k_{\theta_0}/3\})$. Suppose that $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3) = \emptyset$. However, since $i(S^1) \cap (\mathbb{R} \times \{k_{\theta_0}\}) \neq \emptyset$ and $i(S^1) \cap (\mathbb{R} \times \{0\}) \neq \emptyset$, this is inconsistent with the fact that $i(S^1)$ is connected. Suppose that the set $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3)$ has only one point, and put $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3) = \{q\}$. However, since in the circumstance $i(S^1) - \{q\}$ is not connected, this is a contradiction. Hence, we see that the set $\pi_{\theta_0}^{-1}(2k_{\theta_0}/3)$ has two points at least. Let p_1, p_2 be the two points such that

$$p_1 = \left(p_{11}, \frac{2}{3}k_{\theta_0}\right), p_2 = \left(p_{21}, \frac{2}{3}k_{\theta_0}\right) \in \pi_{\theta_0}^{-1}(2k_{\theta_0}/3) \quad (p_{11} < p_{21}).$$

FIGURE 2. $\triangle aa'b'$ and the point b

Then, suppose that there exist two points $b, b' \in i(S^1)$ ($b \neq b'$) such that

$$(1) \quad d_{(p_1, p_2)}|_{i(S^1)}(b) = d_{(p_1, p_2)}|_{i(S^1)}(b'),$$

where $b = (b_1, b_2)$ and $b' = (b'_1, b'_2)$. Note that $b \neq b'$, and by calculating (1) we obtain the following:

$$\begin{aligned} b'_1 &= b_1, \\ b'_2 &= -b_2 + \frac{4}{3}k_{\theta_0}. \end{aligned}$$

Since $b \neq b'$ and $b_1 = b'_1$, we may assume that $b_2 < b'_2$. Now, we show that $0 < b_1 < 1$. If $b_1 > 1$, $b_1 < 0$, we have $d|_{i(S^1) \times i(S^1)}(a, b) > 1$, $d|_{i(S^1) \times i(S^1)}(a', b) > 1$, respectively. These are contradictory with the fact that the maximal value of $d|_{i(S^1) \times i(S^1)}$ is 1. If $b_1 = 0$, then there exist two points b, b' in $\pi_\pi^{-1}(0)$ and if $b_1 = 1$, then there exist two points b, b' in $\pi_0^{-1}(1)$. Since the maximal value of π_π is 0 and the maximal value of π_0 is 1, these are inconsistent with (I). Hence, we have $0 < b_1 < 1$. Since $b_2 < 4k_{\theta_0}/3$, $b'_2 < 4k_{\theta_0}/3$, by seeing (2) we get $b'_2 > 0$, $b_2 > 0$, respectively.

Put

$$\triangle aa'b' = \{t_1 a + t_2 a' + t_3 b' \in \mathbb{R}^2 \mid t_1, t_2, t_3 \geq 0, t_1 + t_2 + t_3 = 1\}.$$

Since the set $i(S^1)$ has three points a, a', b' and by Lemma 4.4 $i(S^1) = \partial \text{conv}(i(S^1))$, it is clearly seen that $\triangle aa'b' \subset \text{conv}(i(S^1))$. Then, we have $(\triangle aa'b')^\circ \subset (\text{conv}(i(S^1)))^\circ$, where $A^\circ \subset \mathbb{R}^2$ is a set which consists of the interior points of A . It is not hard to see that $b \in (\triangle aa'b')^\circ$. Hence, we have $b \in (\text{conv}(i(S^1)))^\circ$. However, this is inconsistent with $b \in \partial \text{conv}(i(S^1))$.

Next, in the circumstance of (II), we shall prove that $d_{(p_1, p_2)}|_{i(S^1)}$ is injective. Then, there exists $\theta_1 \in [0, 2\pi)$ such that the set $\pi_{\theta_1}^{-1}(k_{\theta_1})$ has two points at least, where k_{θ_1} is the maximal value of π_{θ_1} . Let p_1, p_2 be two points in $\pi_{\theta_1}^{-1}(k_{\theta_1})$. By transforming the origin and basis of the linear space $\mathbb{R}^2$ if necessary, we may assume from the first that $p_1 = (0, 0)$, $p_2 = (1, 0)$, and $i(S^1) \subset \mathbb{R} \times \{x \in \mathbb{R} \mid x \leq 0\}$. Then, note that $\theta_1 = \pi/2$, $k_{\theta_1} = 0$. Now, suppose that there exist $c = (c_1, c_2)$, $c' = (c'_1, c'_2) \in i(S^1)$ ($c \neq c'$) such that

$$(2) \quad d_{(p_1, p_2)}|_{i(S^1)}(c) = d_{(p_1, p_2)}|_{i(S^1)}(c').$$

Note that $c \neq c'$, and by calculating (2), we obtain

$$\begin{aligned} c_1 &= c'_1, \\ c_2 &= -c'_2. \end{aligned}$$

Since $c_2 \leq 0$ and $c'_2 \leq 0$, it follows that $c_2 = c'_2 = 0$. However, this contradicts to the assumption $c \neq c'$. $\square$

ACKNOWLEDGEMENT

The authors would like to thank O. Saeki and M. Nishioka for their valuable comments.

REFERENCES

- [1] Morris W. Hirsch, Differential Topology, Springer-Verlag, New York Heidelberg Berlin, 1976.
- [2] J. N. Mather, Generic projections, Ann. of Math. (2) 98 (1973), 226-245.
- [3] J. W. Bruce and N. P. Kirk, Generic projections of stable mappings, Bull. London Math. Soc. 32 (2000), no. 6, 718-728.
- [4] M. Berger, Geometry I, Springer-Verlag, Berlin, 2009.

GRADUATE SCHOOL OF ENVIRONMENT AND INFORMATION SCIENCES, YOKOHAMA NATIONAL UNIVERSITY, YOKOHAMA 240-8501, JAPAN.

E-mail address: 19881126@george24.com

RESEARCH GROUP OF MATHEMATICAL SCIENCES, RESEARCH INSTITUTE OF ENVIRONMENT AND INFORMATION SCIENCES, YOKOHAMA NATIONAL UNIVERSITY, YOKOHAMA 240-8501, JAPAN.

E-mail address: nishimura-takashi-yx@ynu.jp